

ON INFINITELY COHOMOLOGOUS TO ZERO OBSERVABLES

AMANDA DE LIMA AND DANIEL SMANIA

ABSTRACT. We show that for a large class of piecewise expanding maps T , the bounded p -variation observables u_0 that admits an infinite sequence of bounded p -variation observables u_i satisfying

$$u_i = u_{i+1} \circ T - u_{i+1}$$

are constant. The method of the proof consists in to find a suitable Hilbert basis for $L^2(hm)$, where hm is the unique absolutely continuous invariant probability of T . In terms of this basis, the action of the Perron-Frobenius and the Koopman operator on $L^2(hm)$ can be easily understood. This result generalizes earlier results by Bamón, Kiwi, Rivera-Letelier and Urzúa in the case $T(x) = \ell x \bmod 1$, $\ell \in \mathbb{N} \setminus \{0, 1\}$ and Lipschitzian observables u_0 .

1. INTRODUCTION

Let $T: I \rightarrow I$ be a dynamical system. Consider the *cohomological operator* defined by

$$\mathcal{L}(\psi) = \psi \circ T - \psi,$$

Given an observable, that is, a function $u_0: I \rightarrow \mathbb{R}$, one can ask if there exists a solution u_1 to the *Livsic cohomological equation*

$$\mathcal{L}(u_1) = u_0.$$

Such equation was intensively studied after its introduction by the seminal work of Livsic. These studies mainly concerns to the existence and regularity of the solution u_1 .

Let μ to be an invariant probability measure of T . We say that a function $u: I \rightarrow \mathbb{R}$ in $L^1(\mu)$ is cohomologous to zero if there is a function $w: I \rightarrow \mathbb{R}$ in $L^1(\mu)$ such that

$$u = \mathcal{L}(w).$$

A observable u_0 is *infinitely cohomologous to zero* if there exists a sequence of functions $u_n \in L^1(\mu)$, $n \in \mathbb{N}$, such that $\mathcal{L}^n u_n = u_0$, for all $n \in \mathbb{N}$.

In [3], Bamón, Kiwi, Rivera-Letelier and Urzúa consider the expanding maps defined by

$$T_\ell(x) = \ell x \bmod 1,$$

where $\ell \geq 2$ is an integer. The Lebesgue measure on $[0, 1]$ is invariant by T_ℓ . They show that every non-constant Lipschitzian function $u: I \rightarrow \mathbb{R}$ is not infinitely

Date: July 28, 2009.

2000 Mathematics Subject Classification. 37C30, 37E05, 37A05.

Key words and phrases. cohomological equation, Livsic cocycle, exact measure, bounded variation, piecewise expanding map.

A. de L. is partially supported by FAPESP 04/12117-0. D.S. is partially supported by CNPq 470957/2006-9 and 310964/2006-7, FAPESP 2003/03107-9 and 2008/02841-4.

cohomologous to zero. In this work we generalize this result to a much larger class of observables and piecewise expanding maps.

Let I be an interval. We say that $T: I \rightarrow I$ is a *piecewise monotone* map if there exists a partition by intervals $\{I_1, \dots, I_m\}$ of I such that for each $i \leq m$ the map T is continuous and strictly monotone in I_i . A piecewise monotone map is *onto* if furthermore $T(I_i) = I$ for every i . A piecewise monotone map is called *expanding* if T is differentiable on each I_i and

$$\inf_i \inf_{x \in I_i} |T'(x)| > 1.$$

In this work, we will consider mainly maps $T: I \rightarrow I$ satisfying the following conditions:

- (D1) T is piecewise monotone, Lipschitz on each interval of the partition I_i , $i \leq m$. In particular T' is defined almost everywhere and it is an essentially bounded function. We also assume
- (1)
$$\operatorname{ess\,inf}_m |T'| > 0.$$
- (D2) There exists a closed interval $I^* \subset I$ such that for $T(I^*) = I^*$, the orbit of every point of I eventually enters I^* and moreover for every interval $H \subset I$ there is a finite collection of pairwise disjoint open subintervals $H_1, \dots, H_k \subset H$ and n such that T^n is a homeomorphism on H_i and

$$\operatorname{int} I^* \subset \cup_i T^n(H_i).$$

- (D3) T has a horseshoe, that is, there are three open intervals $J_1, J_2 \subset J \subset I^*$, with $J_1 \cap J_2 = \emptyset$, such that T is a homeomorphism on each J_i and $T(J_i) = J$, $i = 1, 2$.
- (D4) T has an invariant probability μ that is absolutely continuous with respect to the Lebesgue measure m , so

$$\mu(A) = \int_A h \, dm$$

for some $h \in L^1(m)$. We will denote $\mu = hm$, where $h \in L^1(m)$ and m is the Lebesgue measure on I . Moreover μ is exact and there exist a, b such that

$$(2) \quad 0 < a \leq h(x) \leq b < \infty$$

for hm -almost every x and the support of μ is I^* .

Our main result is:

Theorem 1. *Let T satisfying D1-D4 and let $u_0: I^* \rightarrow \mathbb{R}$ be an observable with bounded p -variation. Then either u_0 is constant hm -almost everywhere or there exist $m \geq 0$ and bounded p -variation functions $u_i: I^* \rightarrow \mathbb{R}$, with $u_i \in L^1(hm)$, with $i \leq m$, which are unique up to an addition by a constant, such that*

- We have

$$\mathcal{L}^i u_i = u_0 \text{ in } L^1(hm),$$

for every $i \leq m$.

- For every function ρ with bounded p -variation and every $c \in \mathbb{R}$ we have $\mathcal{L}\rho \neq u_m + c$ on $L^1(hm)$.

We can improve this result with additional assumptions on T and u_0 which are satisfied in many interesting situations.

Theorem 2. *Let T be a transformation satisfying D1-D4 and suppose that the Banach space of functions $\mathbb{B}$ and T satisfies D5. Let $u_0 \in \mathbb{B}$ be an observable. Then either u_0 is constant hm -almost everywhere or there exist a unique $m \geq 0$ and functions $u_i : I^* \rightarrow \mathbb{R}$, with $u_i \in L^1(hm)$, with $i \leq m$, which are unique up to an addition by a constant, such that*

- We have

$$\mathcal{L}^i u_i = u_0 \text{ in } L^1(hm)$$

for every $i \leq m$.

- For every function $\rho \in L^1(hm)$ and every $c \in \mathbb{R}$ we have $\mathcal{L}\rho \neq u_m + c$ on $L^1(hm)$

Moreover u_i belongs to $\mathbb{B}$, for $i \leq m$.

We postpone the definition of condition D5 to Section 5, since it is a little more technical.

Remark 1.1. There are plenty of examples of transformations $T : I \rightarrow I$ satisfying D1-D4. Let T be a piecewise monotone, expanding map, C^2 on each I_i . Consider the $m \times m$ matrix $A_T = (a_{ij})$ defined by $a_{ij} = 1$ if

$$\overline{T(\text{int } I_i)} \subset \text{int } I_j,$$

and $a_{ij} = 0$ otherwise. Here the closure and interior are taken with respect to the topology of $[0, 1]$. Suppose that $A_T^k > 0$ for some k . Then T satisfies D1, D2 and D4 and some iteration of T satisfies D1-D4, with $I^* = I$. If we add the assumption that T has a horseshoe, then T fulfills D1-D4. The space of bounded variation functions $BV(I^*)$ and T satisfies D5.

Remark 1.2. A class of examples satisfying D1-D4 are β -transformations $T(x) = \beta x \bmod 1$, with $\beta \geq 2$, $\beta \in \mathbb{R}$, $I^* = I$. The space of bounded variation functions $BV(I^*)$ and T satisfies D5.

Remark 1.3. Let $T : [-1, 1] \rightarrow [-1, 1]$ be a continuous map with $T(-1) = T(1) = -1$, C^2 on the intervals $[-1, 0]$ and $[0, 1]$, with $T' > 0$ in $[-1, 0]$ and $T' < 0$ in $[0, 1]$ and $T(-x) = T(x)$ for every $x \in [-1, 1]$. Define

$$\theta = \inf_x |T'(x)|.$$

If $\theta > 1$ then there exists a unique fixed point $p \in [0, 1]$. Define $J = [-p, p]$. If $\theta > \sqrt{2}$ then T^2 has a horseshoe in J and satisfies D1-D4 with $I^* = [T^2(0), T(0)]$. The space of bounded variation functions $BV(I^*)$ and T satisfies D5.

Remark 1.4. Let $T : I \rightarrow I$ be a piecewise expanding and onto map, $C^{1+\alpha_0}$ in each I_i , $\alpha_0 \in (0, 1)$. Then T satisfies D1 – D4, with $I^* = I$. The space of Holder continuous functions $C^\alpha(I)$, for $\alpha \leq \alpha_0$, and T satisfies D5.

Remark 1.5. Let $T : I \rightarrow I$ be a piecewise expanding map, linear in each I_i . Suppose that T has a horseshoe and satisfies the conditions on the matrix A_T as in Remark 1.1. One can prove using the results of Wong [6] that T satisfies D1 – D4, with $I^* = I$. The space of bounded p -variation functions $BV_p(I)$, with $p \geq 1$, and T satisfies D5.

1.1. Topological Results. Replacing Lipschitzian by bounded p -variation observables has the advantage to allow us to obtain results similar to Theorems 1 and 2 to maps which are just *topologically conjugate* with maps satisfying the assumptions of those theorems.

We will say that two functions $f, g: W \rightarrow \mathbb{R}$ are equal except in a countable subset, $f = g$ on W (e.c.s.) if $\{x \in W: f(x) \neq g(x)\}$ is countable.

Theorem 3. *Let $H: I \rightarrow I$ be a homeomorphism, let T be a piecewise monotone map and $\tilde{T}$ satisfying D1-D4. Suppose that*

$$H \circ \tilde{T} = T \circ H$$

in I (e.c.s.). Let $u_0: H(I^*) \rightarrow \mathbb{R}$ be an observable with bounded p -variation. Then either u_0 is constant in $H(I^*)$ (e.c.s.) or there exist an unique $m \geq 0$ and bounded p -variation functions $u_i: H(I^*) \rightarrow \mathbb{R}$, with $i \leq m$, which are unique up to an addition by a constant (e.c.s.), such that

- *We have*

$$\mathcal{L}^i u_i = u_0,$$

on $H(I^*)$ (e.c.s.) for every $i \leq m$.

- *For every function ρ with bounded p -variation and every $c \in \mathbb{R}$ we have $\mathcal{L}\rho \neq u + c$ in a non-empty open subset in $H(I^*)$.*

Theorem 4. *Let $H: I \rightarrow I$ be a homeomorphism, let T be a piecewise monotone map and $\tilde{T}$ satisfying D1-D4. Suppose that*

$$H \circ \tilde{T} = T \circ H$$

in I (e.c.s.). Suppose that the space of functions with bounded p_0 -variation BV_{p_0, I^*} and $\tilde{T}$ satisfies D5. Let $u_0: H(I^*) \rightarrow \mathbb{R}$ be an observable with bounded p_0 -variation. Then either u_0 is constant in $H(I^*)$ (e.c.s.) or there exist an unique $m \geq 0$ and bounded borelian functions $u_i: H(I^*) \rightarrow \mathbb{R}$, with $i \leq m$, which are unique up to an addition by a constant (e.c.s.), such that

- *We have*

$$\mathcal{L}^i u_i = u_0,$$

on $H(I^*)$ (e.c.s.) for every $i \leq m$.

- *We have $\mathcal{L}\rho \neq u + c$*
 - A. *in an uncountable subset of $H(I^*)$, if ρ is a Borel measurable, bounded function and $c \in \mathbb{R}$.*
 - B. *in a non-empty open subset of $H(I^*)$, if ρ is a Borel measurable, bounded function which is continuous in $H(I^*)$ (e.c.s.) and $c \in \mathbb{R}$.*

Moreover u_i has bounded p_0 -variation, $i \leq m$.

Remark 1.6. Let $T: [0, 2] \rightarrow [0, 2]$ be a piecewise monotone, onto map, C^1 in $[0, 1]$ and $[1, 2]$, with $T(0) = 0$, $T' \geq \lambda > 1$ in $[1, 2]$ and $T'(x) > 1$ in $x \in (0, 1)$ and $T'(0) = 1$. Then T is conjugate with $\tilde{T}(x) = 2x \bmod 1$, so T satisfies the assumptions of Theorems 3 and 4, considering $p_0 = 1$ in Theorem 4.

Remark 1.7. Let $T: [-1, 1] \rightarrow [-1, 1]$, $T(-1) = T(1) = -1$, C^3 in $[-1, 1]$, $T'(0) = 0$, $T' > 0$ on $[-1, 0)$, $T' < 0$ on $(0, 1]$. If T has negative Schwarzian derivative and non-renormalizable then T is conjugate with a tent map $\tilde{T}_\beta: [-1, 1] \rightarrow [-1, 1]$, defined as $\tilde{T}(x) = -\beta|x| + \beta - 1$, with $\beta = \exp(h_{top}(T))$. Here $h_{top}(T)$ denotes the

topological entropy of T . If $h_{top}(T) \geq \ln(2)/2$ then T^2 satisfies the assumptions of Theorems 3 and 4, considering $p_0 = 1$ in Theorem 4.

2. PRELIMINARIES

In this section we present some notations and definitions.

Definition 2.1. Given a function $f : I \rightarrow \mathbb{C}$ and $p \geq 1$, we define the p -variation of f by

$$v_{p,I}(f) = \sup \left(\sum_{i=1}^n |g(a_i) - g(a_{i-1})|^p \right)^{\frac{1}{p}},$$

where the supremum is taken over all finite sequences $a_0 < a_1 < \dots < a_n$, $a_i \in I$.

We say that f has bounded p -variation if

$$v_{p,I}(f) < \infty.$$

It is convenient to us to identify functions u and v defined on I so that $u = v$ up to a countable subset of I . We write $u \sim v$. The set of equivalence classes $[f]$ with respect to the relation $\sim$ such that

$$v_{p,I}([f]) = \inf_{f \sim g} v_{p,I}(g) < \infty$$

will be called the space of the functions on I with bounded p -variation and denoted $BV_{p,I}$. The function $f \rightarrow v_{p,I}([f])$ is a pseudo-norm on $BV_{p,I}$. We can define a norm by

$$|[f]|_{BV_{p,I}} = \inf_{g \sim f} (\sup |g| + v_{p,I}(g)).$$

$(BV_{p,I}, |\cdot|_{BV_{p,I}})$ is a Banach space. As usual, from now on we will omit the brackets $[\cdot]$ in the notation of the equivalence classes.

Note that $1/p$ -Hölder continuous functions have bounded p -variation. When $p = 1$, we say that the function has bounded variation.

Remark 2.2. One of the greatest advantages of dealing with p -bounded variation observables, in opposition to either Hölder or Lipschitzian ones, for instance, is that the pseudo-norm $v_{p,I}$ is invariant by homeomorphisms, that is, if $h : J \rightarrow I$ is a homeomorphism and $f : I \rightarrow \mathbb{R}$ is an observable then

$$v_{p,I}(f) = v_{p,J}(f \circ h).$$

Definition 2.3. Given a piecewise monotone, expanding map T , define the Perron-Frobenius operator associated to T by

$$\Phi_T f(x) = \sum_{j \in J} f(\sigma_j(x)) \frac{1}{|T'(\sigma_j x)|} \mathbb{1}_{T(I_j)}(x),$$

where $\sigma_j : T(I_j) \rightarrow I_j$ stands for the inverse branch of T restricted to I_j and $\mathbb{1}_J$ denotes the characteristic function of the set J .

The main properties of Φ_T are:

i) Φ_T is a continuous linear operator on $L^1(hm)$.

ii) $\int_0^1 \Phi_T f \cdot g \, dm = \int_0^1 f \cdot g \circ T \, dm$, where $f \in L^1(m)$ and $g \in L^\infty(m)$.

iii) $\Phi_T f = f$ if and only if the measure $\mu = fm$ is invariant by T .

3. A SPECIAL BASIS OF $L^2(hm)$

Consider the Hilbert space $L^2(hm)$ with the inner product

$$\langle u, w \rangle_{hm} = \int uwh \, dm.$$

Indeed $\langle u, w \rangle_{hm}$ is well defined even for $u \in L^k(hm)$ and $w \in L^b(hm)$, with $k, b \in [1, \infty) \cup \{+\infty\}$ satisfying

$$\frac{1}{k} + \frac{1}{b} = 1.$$

In this section we will build a Hilbert basis for $L^2(hm)$. Consider the bounded linear operator $P : L^k(hm) \rightarrow L^k(hm)$, $k \geq 1$, defined by

$$P(u) = \frac{\Phi(uh)}{h}.$$

Due Eq. (2), the operator P is well defined. Let $\mathcal{B} = \{\varphi_i\}_{i \in \mathbb{N}}$ be an orthonormal basis for

$$\text{Ker}(P) = \{u \in L^2(hm) \text{ s.t. } P(u) = 0\}.$$

Define

$$\mathcal{W} = \{\varphi_i \circ T^j : \varphi_i \in \mathcal{B} \text{ e } j \in \mathbb{N}\} \cup \{\mathbb{1}_{I^*}\}.$$

where $\mathbb{1}$ denotes the constant function $\mathbb{1}(x) = 1$.

Proposition 3.1. *$\mathcal{W}$ is a Hilbert basis for $L^2(hm)$. Indeed we can choose $\mathcal{B}$ such that $\mathcal{W} \subset L^\infty(hm)$.*

Remark 3.2. A very interesting example of this theorem is given by the function $T : [0, 1] \rightarrow [0, 1]$ defined by $T(x) = \ell x \bmod 1$, with $\ell \in \mathbb{N} \setminus \{0, 1\}$. In this case the Ruelle-Perron-Frobenius operator is just

$$(\Phi_T \psi)(x) = \frac{1}{\ell} \sum_{i=0}^{\ell-1} \psi\left(\frac{x+i}{\ell}\right).$$

The Lebesgue measure m is an invariant probability, so $P = \Phi_T$. Moreover

$$\mathcal{B} = \{\sin(2\pi n x), \cos(2\pi n x) : \ell \text{ does not divide } n\}$$

is a basis for $\text{Ker } P$. Note that

$$\sin(2\pi n T^j(x)) = \sin(2\pi n \ell^j x) \text{ and } \cos(2\pi n T^j(x)) = \cos(2\pi n \ell^j x),$$

so the corresponding set $\mathcal{W}$ is just the classical Fourier basis of $L^2([0, 1])$.

It is easy to see that the Koopman operator $U : L^k(hm) \rightarrow L^k(hm)$, $k \geq 1$, defined by

$$U(w) = w \circ T,$$

is the adjoint operator of P , that is

$$(3) \quad \langle P(u), w \rangle_{hm} = \langle u, U(w) \rangle_{hm}$$

for every $u \in L^k(hm)$ and $w \in L^b(hm)$. Moreover

$$P \circ U(f) = f$$

for every $f \in L^1(hm)$.

Lemma 3.3. $\mathcal{W}$ is an orthonormal set.

Proof. Indeed

$$\begin{aligned} |\mathbb{1}_{I^*}|_{L^2(hm)} &= 1, \\ |\varphi_i \circ T^j|_{L^2(hm)}^2 &= |\varphi_i|_{L^2(hm)}^2 = 1. \end{aligned}$$

Futhermore if

$$(i_1, j_1) \neq (i_2, j_2)$$

then either $j_1 = j_2$, so we have

$$\langle \varphi_{i_1} \circ T^{j_1}, \varphi_{i_2} \circ T^{j_2} \rangle_{hm} = \langle \varphi_{i_1}, \varphi_{i_2} \rangle_{hm} = 0,$$

or without loss of generality we can assume $j_1 < j_2$ and

$$\langle \varphi_{i_1} \circ T^{j_1}, \varphi_{i_2} \circ T^{j_2} \rangle_{hm} = \langle \varphi_{i_1}, \varphi_{i_2} \circ T^{j_2-j_1} \rangle_{hm} = \langle P^{j_2-j_1}(\varphi_{i_1}), \varphi_{i_2} \rangle_{hm} = 0,$$

and

$$\langle \varphi_{i_1} \circ T^{j_1}, \mathbb{1}_{I^*} \rangle_{hm} = \int \varphi_i \circ T^j h \, dm = \int \varphi_i h \, dm = \int P(\varphi_i) h \, dm = 0.$$

□

Lemma 3.4. *There exists a countable set of functions $\Lambda \subset L^\infty(hm) \cap \text{Ker}(P)$ with the following property: Let $w \in L^k(hm)$, with $k \geq 1$. If for all $\varphi \in \Lambda$ we have*

$$\int w \varphi h \, dm = 0,$$

then there exists $\beta \in L^k(hm)$ such that

$$w = \beta \circ T$$

hm-almost everywhere. In particular $\text{Ker}(P)^\perp = U(L^2(hm))$.

Proof. Note that for the existence of $\beta \in L^k(hm)$ such that $w = \beta \circ T$ is necessary and sufficient that for hm -almost every $y \in I$ we have

$$(4) \quad \sharp\{w(x) : h(x) \neq 0 \text{ and } T(x) = y\} = 1.$$

Indeed, if the Eq. (4) holds then for every y satisfying (4), choosing x such that $T(x) = y$ and $h(x) \neq 0$ we can define

$$\beta(y) = w(x).$$

If y does not satisfy (4), define $\beta(y) = 0$. Of course $w = \beta \circ T$ hm -almost everywhere and, since hm is an invariant measure of T , β belongs to $L^k(hm)$.

Let C_i be the set of points $x_0 \in I$ such that the function

$$(5) \quad F_i(a) = \int_0^a w \circ \sigma_i(Tx) \cdot \mathbb{1}_{T(I_i)}(T(x)) \cdot h(x) dm(x)$$

has derivative $w \circ \sigma_i(T(x_0)) \mathbb{1}_{T(I_i)}(T(x_0)) h(x_0)$ at $a = x_0$. The function in the above integral belongs to $L^1(m)$, so by the Lebesgue differentiation theorem the set

$$C = \cap_i C_i \setminus \cup_i \partial I_i$$

has full Lebesgue measure in I . Since T is piecewise Lipschitz we obtain that

$$m(T(I \setminus C)) = 0.$$

Suppose that Eq. (4) does not hold for hm -almost every $y \in I^*$. Then it is not true that Eq. (4) holds for hm -almost every $y \in I \setminus T(I \setminus C)$. Since hm -almost

every point has at least one preimage x with $h(x) \neq 0$, we conclude that there exists $y_0 \in I \setminus T(I \setminus C)$ and two inverse branches of T , denoted by σ_1 and σ_2 such that y_0 belongs to the interior of $T(I_1) \cap T(I_2)$ and furthermore

$$w \circ \sigma_1(y_0) \neq w \circ \sigma_2(y_0), h(\sigma_1(y_0)) \neq 0, h(\sigma_2(y_0)) \neq 0.$$

We can assume

$$w \circ \sigma_1(y_0) > w \circ \sigma_2(y_0),$$

so

$$(6) \quad w \circ \sigma_1 \circ T \circ \sigma_2(y_0) \mathbb{1}_{T(I_1)} \circ T \circ \sigma_2(y_0) h \circ \sigma_2(y_0) > w \circ \sigma_2 \circ T \circ \sigma_2(y_0) \mathbb{1}_{T(I_2)} \circ T \circ \sigma_2(y_0) h \circ \sigma_2(y_0).$$

Since $\sigma_2(y_0) \in C$, the derivatives of the functions F_1 and F_2 at $a = \sigma_2(y_0)$ are the left and right hand sides of Eq. (6) respectively, so there exists $\varepsilon > 0$ such that for every closed non degenerate interval $\tilde{I}_2$ satisfying

$$(7) \quad \sigma_2(y_0) \in \tilde{I}_2 \subset (\sigma_2(y_0) - \varepsilon, \sigma_2(y_0) + \varepsilon) \cap I_2$$

we have

$$\int_{\tilde{I}_2} w \circ \sigma_1(Tx) \mathbb{1}_{T(I_1)} \circ T(x) \cdot h(x) dm(x) > \int_{\tilde{I}_2} w \circ \sigma_2(Tx) \mathbb{1}_{T(I_2)} \circ T(x) \cdot h(x) dm(x).$$

Choose an interval $\tilde{I}_2$ satisfying Eq. (7) and small enough such that $T(\tilde{I}_2) \subset T(I_1)$. We can assume without loss of generality that $\partial \tilde{I}_2 \subset \mathbb{Q}$. Then

$$\int_{\tilde{I}_2} w \circ \sigma_1(Tx) \cdot h(x) dm(x) > \int_{\tilde{I}_2} w \circ \sigma_2(Tx) \cdot h(x) dm(x).$$

Let $\tilde{I}_1 := \sigma_1(T(\tilde{I}_2)) \subset I_1$. Define φ as

$$(8) \quad \varphi(x) = \begin{cases} -\frac{|T'(x)|}{|T'(\sigma_2(Tx))|} \cdot \frac{h(\sigma_2(Tx))}{h(x)} & \text{if } x \in \tilde{I}_1, \\ 1 & \text{if } x \in \tilde{I}_2, \\ 0 & \text{otherwise.} \end{cases}$$

Note that $\varphi \in L^\infty(hm)$ and $\Phi(\varphi h) = 0$.

Hence

$$\begin{aligned} \int w \varphi h dm &= \\ &= \int_{\tilde{I}_1} w \varphi h dm + \int_{\tilde{I}_2} w \varphi h dm \\ &= \int_{\tilde{I}_1} w \cdot \left(-\frac{|T'|}{|T' \circ \sigma_2 \circ T|} \frac{h \circ \sigma_2 \circ T}{h} \right) h dm + \int_{\tilde{I}_2} w h dm. \end{aligned}$$

Since $\sigma_2 \circ T : \tilde{I}_1 \rightarrow \tilde{I}_2$ is Lipschitzian and monotone increasing, we can make a change of variables to get

$$\begin{aligned} & - \int_{\tilde{I}_1} w \frac{|T'|}{|T' \circ \sigma_2 \circ T|} \frac{h \circ \sigma_2 \circ T}{h} h \, dm + \int_{\tilde{I}_2} w h \, dm \\ &= - \int_{\tilde{I}_2} w \circ \sigma_1 \circ T \cdot h \, dm + \int_{\tilde{I}_2} w \circ \sigma_2 \circ T \cdot h \, dm \\ &= - \int_{\tilde{I}_2} w \circ \sigma_1 \circ T \cdot h \, dm + \int_{\tilde{I}_2} w \circ \sigma_2 \circ T \cdot h \, dm \\ &< - \int_{\tilde{I}_2} w \circ \sigma_2 \circ T \cdot h \, dm + \int_{\tilde{I}_2} w \circ \sigma_2 \circ T \cdot h \, dm = 0. \end{aligned}$$

Therefore

$$\int w \varphi h \, dm \neq 0.$$

Let Λ be the set of functions φ of the form in Eq. (8), with

- The intervals $\tilde{I}_j \subset I_{i_j}$, $j = 1, 2$, and $\sigma_2 : T(I_{i_2}) \rightarrow I_{i_2}$ is the inverse of $T : I_{i_2} \rightarrow T(I_{i_2})$.
- $T(\tilde{I}_2) = T(\tilde{I}_1)$.
- $\partial \tilde{I}_2 \subset \mathbb{Q}$.

Then it is easy to see that Λ is countable and $\Lambda \subset L^\infty(hm) \cap \text{Ker} P$ and, by the argument above, Λ has the wished property.

In particular for $k = 2$ we obtain $\text{Ker}(P)^\perp \subset U(L^2(hm))$. The inclusion $U(L^2(hm)) \subset \text{Ker}(P)^\perp$ follows from Eq. (3). $\square$

Proposition 3.5. *Let Λ be as in Lemma 3.4. Let $u : I \rightarrow \mathbb{R}$ be a non constant function in $L^1(hm)$. Then there exists $\varphi \in \Lambda$, and an integer $p \geq 0$ such that*

$$\int u \varphi \circ T^j \cdot h \, dm = 0, \text{ for all } 0 \leq j < p$$

and

$$\int u \varphi \circ T^p \cdot h \, dm \neq 0.$$

Proof. Suppose that, for all $\varphi \in \Lambda$ and for all $k \geq 0$

$$(9) \quad \int u \varphi \circ T^k \cdot h \, dm = 0.$$

We claim that for every n there exists $\beta_n \in L^1(hm)$ such that

$$(10) \quad u = \beta_n \circ T^n.$$

Indeed, choosing $k = 0$ in Eq. (9) we obtain that for all $\varphi \in \Lambda$

$$\int u \varphi h \, dm = 0.$$

By Lemma 3.4, there exists $\beta_1 \in L^1(hm)$ such that

$$u = \beta_1 \circ T.$$

Suppose by induction that $u = \beta_n \circ T^n$, with $\beta_n \in L^1(hm)$. By Eq. (9) when $k = n$, for all $\varphi \in \Lambda$ we have

$$\int \beta_n \varphi h \, dm = \int \beta_n \circ T^n \cdot \varphi \circ T^n \cdot h \, dm = \int u \varphi \circ T^n \cdot h \, dm = 0.$$

Hence, by Lemma 3.4, there exists $\beta_{n+1} \in L^1(hm)$ such that

$$\beta_n = \beta_{n+1} \circ T \Rightarrow u = \beta_{n+1} \circ T^{n+1}.$$

Since the measure hm is an exact measure, we can conclude that u is a constant function. So $u = 0$. □

Corollary 3.6. *Let $u : I \rightarrow \mathbb{R}$ be a non constant function in $L^2(hm)$. Then there exist $\varphi_i \in \mathcal{B}$ and an integer $p \geq 0$ such that*

$$\langle u, \varphi_i \circ T^j \rangle_{hm} = 0 \text{ for all } 0 \leq j < p$$

and

$$\langle u, \varphi_i \circ T^p \rangle_{hm} \neq 0.$$

Proof. Suppose that for every $\varphi_i \in \mathcal{B}$ and every $j \in \mathbb{N}$

$$\langle u, \varphi_i \circ T^j \rangle_{hm} = 0.$$

Since $\mathcal{B}$ is a base for $\text{Ker}(P)$ and $U^j : L^2(hm) \rightarrow L^2(hm)$ is an isometry, it follows that

$$\int \varphi \circ T^j \cdot u \cdot h \, dm = 0$$

for every $\varphi \in \text{Ker}(P)$ and $j \in \mathbb{N}$. This contradicts Proposition 3.5. □

Proof of Proposition 3.1. It follows from Lemma 3.3 and Corollary 3.6 that $\mathcal{B}$ is a basis of $L^2(hm)$. To construct a basis $\mathcal{W} \subset L^\infty(hm)$, consider an enumeration of the set $\Lambda = \{\psi_i\}$ defined in Lemma 3.4. Apply the Gram-Schmidt process in the sequence ψ_i to obtain a sequence $\tilde{\psi}_i$ of two by two orthogonal functions. Discarding the null functions and normalizing the remaining functions, we obtain an orthonormal set of functions $\hat{\mathcal{B}}$. Due Lemma 3.4

$$\overline{\text{span}(\hat{\mathcal{B}})} = \overline{\text{span}(\Lambda)} = \text{Ker } P,$$

so $\hat{\mathcal{B}}$ is a basis of $\text{Ker } P$, so

$$\hat{\mathcal{W}} = \{\phi \circ T^j : \phi \in \hat{\mathcal{B}}, j \in \mathbb{N}\} \cup \{\mathbb{1}_{I^*}\}$$

is a basis of $L^2(hm)$. □

From now on we assume $\mathcal{W} \subset L^\infty(hm)$. Let $u \in L^1(hm)$ and consider the Fourier coefficients of u with respect to the basis $\mathcal{W}$

$$c_{i,j}(u) = \int u \cdot \varphi_i \circ T^j \cdot h \, dm.$$

Proposition 3.7. *The functionals $c_{i,j}$ have the following properties:*

- (1) $c_{i,j}$ is linear on $L^1(hm)$
- (2) $c_{i,j}(U(u)) = c_{i,j-1}(u)$.
- (3) $c_{i,0}(U(u)) = 0$.
- (4) $c_{i,j}(P(u)) = c_{i,j+1}(u)$.

Proof. (1) The proof is straightforward.

(2)

$$\begin{aligned} c_{i,j}(u \circ T) &= \int u \circ T \circ \varphi_i \circ T^j \cdot h \, dm = \int u \circ T \varphi_i \circ T^{j-1} \circ T \cdot h \, dm \\ &= \int u \varphi_i \circ T^{j-1} \cdot h \, dm = c_{i,j-1}(u). \end{aligned}$$

(3)

$$c_{i,0}(u \circ T) = \int u \circ T \cdot \varphi_i h \, dm = \int u \Phi(\varphi_i h) \, dm = 0.$$

(4)

$$\begin{aligned} c_{i,j}(Pu) &= \int P(u) \circ T \varphi_i \circ T^j h \, dm = \int \Phi(uh) \varphi_i \circ T^j \, dm \\ &= \int u \varphi_i \circ T^j \circ T \cdot h \, dm = c_{i,j+1}(u). \end{aligned}$$

□

4. PROOF OF THE THEOREM 1

In this section we will study the linear operator

$$\mathcal{L}u = u \circ T - u$$

acting on functions with bounded variation (resp. $1/p$ -Hölder continuous function) $u : I \rightarrow \mathbb{R}$.

First, we will present some properties and then, at the end of this section, we will prove the theorems announced in introduction. The following results are well know.

Lemma 4.1. *Let $\mathcal{L}$ be the linear operator defined above acting on $L^1(hm)$. Then:*

- (1) *If $f \in \text{Im}(\mathcal{L})$, then $\int f h \, dm = 0$.*
- (2) *$\text{Ker}(\mathcal{L}) = \{f \in L^1(hm) : f \text{ is constant } hm\text{-almost everywhere}\}$.*

Corollary 4.2. *Let $u \in L^1(hm)$ and suppose that there exist functions $v, w \in L^1(hm)$ such that*

$$\mathcal{L}^n v = u = \mathcal{L}^n w.$$

Then $v = w + c$ on $L^1(hm)$, for some $c \in \mathbb{R}$. Moreover if v, w have bounded p -variation then $v = w + c$ on $I^*(e.c.s.)$.

Proof. Define $v_i = \mathcal{L}^{n-i} v$, $w_i = \mathcal{L}^{n-i} w$. We will prove by induction on i that $v_i = w_i$, if $i < n$ and $v_n = w_n + c$, for some $c \in \mathbb{R}$. Indeed, for $i = 0$ we have $w_0 = v_0 = u$. Suppose that $v_i = w_i$, $i < n$. Then

$$\mathcal{L}(v_{i+1} - w_{i+1}) = v_i - w_i = 0,$$

so $v_{i+1} - w_{i+1}$ is hm -almost everywhere constant. If $i + 1 = n$ we are done. If $i + 1 < n$ then $\mathcal{L}v_{i+2} = v_{i+1}$ and $\mathcal{L}w_{i+2} = w_{i+1}$, so

$$\int v_{i+1} h \, dm = \int w_{i+1} h \, dm = 0,$$

which implies $c = 0$. Now assume that u, v and w have bounded p -variation. Since the support of hm is I^* and $v = w + c$ hm -almost everywhere, we have the $v = w + c$ on a set $\Lambda \subset I^*$ such that for every non-empty open subset O of I^* we have that

$O \cap \Lambda$ is a dense and uncountable subset of O . Since v and w have just a countable number of discontinuities in I^* , it follows that $v = w + c$ in I^* (e.c.s.). $\square$

Lemma 4.3. *Let J be an open interval as in D3 and $u \in BV_{p,I^*}$. Then*

$$v_{p,J}(\mathcal{L}u) \geq v_{p,J}(u)$$

for every $n \in \mathbb{N}$.

Proof. Let $J_1, J_2 \subset J$ be as in D3. Since T is a homeomorphism on J_1 and J_2 , by Remark 2.2

$$v_{p,J}(u \circ T) \geq v_{p,J_1}(u \circ T) + v_{p,J_2}(u \circ T) = 2v_{p,J}(u),$$

so

$$v_{p,J}(u \circ T - u) \geq v_{p,J}(u \circ T) - v_{p,J}(u) \geq v_{p,J}(u).$$

$\square$

Lemma 4.4. *There exists C with the following property: Let $u_n : I \rightarrow \mathbb{R}$, $n \leq m+1$, be observables with bounded p -variation, $p \geq 1$, such that for every $n \leq m$*

$$u_n = \mathcal{L}u_{n+1}.$$

Then

$$|u_n|_{L^\infty(hm)} \leq v_{p,I^*}(u_n) \leq C v_{p,I^*}(u_0)$$

for every $n \leq m$.

Proof. Let $J \subset I^*$ be an one interval as in D3. By Lemma 4.3

$$(11) \quad v_{p,J}(u_n) \leq v_{p,J}(u_0)$$

for every $n \geq 0$. By D2 there is a finite collection of pairwise disjoint open intervals $H_1, \dots, H_k \subset J$ and j such that T^j is a homeomorphism on each H_i and

$$(12) \quad \text{int } I^* \subset \bigcup_{i=1}^k T^j(H_i).$$

We claim that for every $\ell \leq j$ and n

$$(13) \quad v_{p,T^\ell(H_i)}(u_n) \leq 2^\ell v_{p,J}(u_0)$$

We will prove this by induction on ℓ . Of course since $H_i \subset J$, Eq. (11) implies that for every $i = 1, \dots, k$

$$(14) \quad v_{p,H_i}(u_n) \leq v_{p,J}(u_0),$$

So Eq. (13) holds for $\ell = 0$. Suppose by induction that Eq. (13) holds for $\ell < j$ and every n . Since T is a homeomorphism on $T^\ell(H_i)$ and $u_{n-1} = u_n \circ T - u_n$ we have

$$\begin{aligned} v_{p,T^{\ell+1}(H_i)}(u_n) &= v_{p,T^\ell(H_i)}(u_n \circ T) \leq v_{p,T^\ell(H_i)}(u_n) + v_{p,T^\ell(H_i)}(u_{n-1}) \\ &\leq 2^{\ell+1} v_{p,J}(u_0). \end{aligned}$$

By Eq (12)

$$v_{p,I^*}(u_n) = v_{p,\text{int } I^*}(u_n) \leq \sum_{i=1}^k v_{p,T^j(H_i)}(u_n) \leq k 2^j v_{p,J}(u_0) \leq k 2^j v_{p,I^*}(u_0).$$

Note that since $u_n = u_{n+1} \circ T - u_{n+1}$ it follows that

$$\int u_n h \, dm = 0,$$

Suppose that

$$\text{ess sup}_{hm} u_n = |u_n|_{L^\infty(hm)}.$$

Then

$$\begin{aligned} 0 = \int u_n h \, dm &\geq \text{ess inf } u_n = (\text{ess inf } u_n - \text{ess sup } u_n) + \text{ess sup } u_n \\ &\geq -v_{p,I^*}(u_n) + |u_n|_{L^\infty(hm)}. \end{aligned}$$

so $|u_n|_{L^\infty(hm)} \leq v_{p,I^*}(u_n)$. We can obtain the same conclusion for the case

$$-\text{ess inf}_{hm} u_n = |u_n|_{L^\infty(hm)},$$

replacing u_n by $-u_n$ in the argument above. $\square$

Proof of Theorem 1. Define by induction the (either finite or infinite) sequence $u_n: I^* \rightarrow \mathbb{R}$ of functions in the following way: u_0 is given. If u_n is defined and there exists a function $v: I^* \rightarrow \mathbb{R}$ with bounded p -variation such that $\mathcal{L}v = u_n$ in $L^1(hm)$, then define

$$u_{n+1} = v - \int v h \, dm.$$

Otherwise the sequence ends with u_n . Note that

$$\mathcal{L}^n u_n = u_0.$$

Define

$$m_0 = \sup\{n \in \mathbb{N}: u_n \text{ is defined}\} \in \mathbb{N} \cup \{\infty\}.$$

We will show that $m_0 < \infty$. Let $m \in \mathbb{N}$, $m \leq m_0$. Recall the basis $\mathcal{W}$ defined in Section 3. By Corollary 3.6 if u_0 is not constant there exist i and $q \geq 0$ such that

$$c_{i,j}(u_0) = \int u_0 \varphi_i \circ T^j \cdot h \, dm = 0, \quad \text{for all } 0 \leq j < q$$

and

$$c_{i,q}(u_0) = \int u_0 \varphi_i \circ T^q \cdot h \, dm \neq 0.$$

By Lemma 4.4 we have that $|u_n|_{L^\infty(hm)} \leq C v_{p,I^*}(u_0)$, so

$$|c_{i,k}(u_n)| = \left| \int u_n \varphi_i \circ T^k \cdot h \, dm \right| \leq \int |u_n| |\varphi_i \circ T^k| |h| \, dm \leq M v_{p,I^*}(u_0).$$

Using Lemma 3.7, we can now use an argument quite similar to [3]. Observe that

$$c_{i,l}(u_{n-1}) = c_{i,l}(u_n \circ T - u_n) = c_{i,l}(u_n \circ T) - c_{i,l}(u_n) = c_{i,l-1}(u_n) - c_{i,l}(u_n),$$

for $l \geq 1$.

For $l = 0$,

$$c_{i,0}(u_{n-1}) = c_{i,0}(u_n \circ T - u_n) = c_{i,0}(u_n \circ T) - c_{i,0}(u_n) = -c_{i,0}(u_n),$$

for $0 < n \leq m$.

Therefore, for $0 < n \leq m$

$$(15) \quad c_{i,l}(u_n) = c_{i,l-1}(u_n) - c_{i,l}(u_{n-1}), \quad \text{for } l \geq 1.$$

$$(16) \quad c_{i,0}(u_{n-1}) = -c_{i,0}(u_n).$$

Since $c_{i,j}(u_0) = 0$ for $0 \leq j < q$, by equations (15) and (16), we can conclude that

$$(17) \quad c_{i,j}(u_n) = 0 \text{ for } 0 \leq j < q \text{ and } 0 \leq n \leq m.$$

Now, by equation (15), considering $l = p$, we have

$$c_{i,q}(u_{n-1}) = c_{i,q-1}(u_n) - c_{i,q}(u_n).$$

By equation (17), for every $n \leq m$

$$(18) \quad c_{i,q}(u_{n-1}) = -c_{i,q}(u_n).$$

By equation (18), we conclude that for $n \leq m$

$$c_{i,q}(u_n) = (-1)^n c_{i,q}(u_0).$$

Considering $l = q + 1$ in the equation (15)

$$c_{i,q+1}(u_n) = (-1)^n c_{i,q}(u_0) - c_{i,q+1}(u_{n-1}) \Rightarrow$$

$$(19) \quad c_{i,q}(u_0) = (-1)^n c_{i,q+1}(u_n) + (-1)^n c_{i,q+1}(u_{n-1}).$$

Putting $n = 1, \dots, m$ in Eq. (19) and adding the resulting equations we obtain

$$(20) \quad m \cdot c_{i,q}(u_0) = (-1)^m c_{i,q+1}(u_m) - c_{i,q+1}(u_0).$$

Therefore,

$$\begin{aligned} m &= \frac{-c_{i,q+1}(u_0) + (-1)^m c_{i,q+1}(u_m)}{c_{i,q}(u_0)} \\ &\leq \frac{|c_{i,q+1}(u_0)| + |c_{i,q+1}(u_m)|}{|c_{i,q}(u_0)|} \\ &\leq \frac{|c_{i,q+1}(u_0)| + M v_{p,I}(u_0)}{|c_{i,q}(u_0)|}. \end{aligned}$$

So m_0 is bounded. Note that by Corollary 4.2, if $v_n \in L^1(hm)$ satisfies $\mathcal{L}^n v_n = u_0$ then $v_n = u_n + c$ in $L^1(hm)$, for some $c \in \mathbb{R}$. This proves the uniqueness statements of Theorem 1. $\square$

5. PROOF OF THE THEOREM 2

Let $(\mathbb{B}, |\cdot|_{\mathbb{B}})$ be a Banach space of real-valued, Lebesgue measurable functions defined on I such that

- (D5) (i) T is a piecewise expanding map.
(ii) There exists C and $p_0 \geq 1$ such that

$$|f|_{BV_{p_0,I^*}} \leq C|f|_{\mathbb{B}}$$

for every $f \in \mathbb{B}$. Here $v_{p,I}(f)$ denotes the p -variation of f on I .

(iii) the Perron-Frobenius operator Φ_T of T is a bounded operator on $\mathbb{B}$ and there exists $h \in \mathbb{B}$, $h > 0$, $\lambda \in [0, 1)$ and an operator $\Psi: \mathbb{B} \rightarrow \mathbb{B}$ such that

$$\Phi_T(f) = \int f \cdot dm \cdot h + \Psi(f),$$

with

$$|\Psi^n|_{\mathbb{B}} \leq C\lambda^n,$$

for every $f \in \mathbb{B}$ and $n \in \mathbb{N}$. Moreover and $\Psi(h) = 0$.

(iv) $1/h \in \mathbb{B}$.

(v) The multiplication

$$(f, g) \rightarrow f \cdot g$$

is a bounded bilinear transformation on $\mathbb{B}$.

(vi) The set $\mathbb{B}$ is dense in $L^1(hm)$.

Remark 5.1. Note that D5.i implies that for every $k \geq 1$ we have

$$\mathbb{B} \subset BV_{p_0, I^*} \subset L^\infty(hm) \subset L^k(hm).$$

Indeed for every $v \in \mathbb{B}$ and $k \geq 1$ we have

$$|v|_{L^k(hm)} \leq |v|_{L^\infty(hm)} \leq \tilde{C}|v|_{BV_{p_0, I^*}} \leq C|v|_{\mathbb{B}}.$$

So D5.iii-v implies that

$$\frac{1}{h} \Psi^j(vh)$$

converges exponentially to zero in $L^k(hm)$, $L^\infty(hm)$, BV_{p_0, I^*} and $\mathbb{B}$.

The proof of Theorem 2 is an immediate consequence of Theorem 1 and

Lemma 5.2. *Let T be a transformation satisfying D1-D4 and suppose that $\mathbb{B}$ and T satisfies D5. Let $u \in \mathbb{B}$ and suppose that there exists $v \in L^1(hm)$ such that*

$$u = \mathcal{L}v$$

on I^* . Then v coincides hm -almost everywhere with a function $v_1 \in \mathbb{B}$.

Proof. The method we are going to use here is very well known for specific kinds certain dynamical systems and observables. See for instance [4] for the case of piecewise expanding maps and bounded variation observables. Since

$$u = v \circ T - v,$$

Applying P^j , $j \geq 1$, we get

$$(21) \quad P^j u = P^{j-1} v - P^j v,$$

Putting $j = 1, \dots, n$ in Eq. (21) and adding the resulting equations we obtain

$$v = P^n v + \sum_{j=1}^n P^j u$$

We claim that $|P^j v|_{L^1(hm)} \rightarrow_j 0$. Indeed, due D5.vi for every $\epsilon > 0$ there exists $w \in \mathbb{B}$ such that $\int w h dm = 0$ and $|v - w|_{L^1(hm)} < \epsilon$. Since P is a contraction on $L^1(hm)$, for every j

$$|P^j v - P^j w|_{L^1(hm)} < \epsilon.$$

Due D5 we have that for every $w \in \mathbb{B}$

$$P^j(w) = \frac{1}{h} \Psi^j(wh),$$

with

$$|\Psi^j|_{\mathbb{B}} \leq C\lambda^j,$$

so for j large enough

$$|P^j v|_{L^1(hm)} < 2\epsilon.$$

This proves our claim. In particular

$$v = \sum_{j=1}^{\infty} P^j u,$$

where the convergence of the series is in $L^1(hm)$. On the other hand, by Remark 5.1 this series converges in $L^1(hm)$ and $\mathbb{B}$ to a function $v_1 \in \mathbb{B}$. So $v = v_1$ hm -almost everywhere. $\square$

Remark 5.3. The finiteness result for the family of cohomological operators

$$\mathcal{L}_\lambda(v) = v \circ T - \lambda v,$$

with $\lambda \in (0, 1]$, $T(x) = \ell x \bmod 1$, for integers $\ell \geq 2$ and Lipchitz observables, obtained in [3, Main Lemma, page 225], can also be generalized for maps described in Remarks 1.1, 1.2 and 1.3, replacing Lipchitz observables by bounded variation observables. The methods to achieve this generalization are quite similar to those in [3], so we will not give a full proof here. It is necessary to use Theorem 2, and to replace in their argument the usual Fourier basis by the basis obtained in Section 3 and the compactness of closed balls centered at zero of the space of Lipchitz functions as subsets of the space of continuous functions by Helly's Theorem, that is, the compactness of closed balls centered at zero of the space of bounded variation functions as subsets of $L^1(hm)$.

6. TOPOLOGICAL RESULTS

Proof of Theorem 3. Define $\tilde{u}_0 = u_0 \circ H$. Then $\tilde{u}_0$ has bounded p -variation. By Theorem 1 there exist bounded p -variation functions $\tilde{u}_i$, $i \leq m$, unique up to a constant, such that

$$\tilde{\mathcal{L}}^i \tilde{u}_i = \tilde{u}_0 \text{ on } L^1(hm),$$

and

$$(22) \quad \tilde{\mathcal{L}}\alpha \neq \tilde{u}_m + c \text{ on } L^1(hm),$$

for every bounded p -variation function α . Here $\tilde{\mathcal{L}}v = v \circ \tilde{T} - v$. Since the support of hm is I^* , it follows that $\tilde{\mathcal{L}}^i \tilde{u}_i = \tilde{u}_0$ in $I^*(e.c.s)$. Define $u_i = \tilde{u}_i \circ H^{-1}$. Then u_i has bounded p -variation and

$$\mathcal{L}^i u_i = u_0 \text{ on } H(I^*)(e.c.s).$$

Suppose that there exist a function ρ with bounded p -variation such that $\mathcal{L}\rho = u_m + c(e.c.s)$. Define $\tilde{\rho} = \rho \circ H$. Then $\tilde{\rho}$ has bounded p -variation and $\tilde{\mathcal{L}}\tilde{\rho} = \tilde{u}_m + c$ on $L^1(hm)$. That contradicts Eq. (22). So $\mathcal{L}\rho \neq u_m + c$ in an uncountable subset of I^* . Since the discontinuities of $\mathcal{L}\rho$ and $u_m + c$ are countable, it follows that there is a continuity point $x_0 \in I^*$ of both functions such that $(\mathcal{L}\rho)(x_0) \neq u_m(x_0) + c$. So there is a non-empty open subset of I^* such that $\mathcal{L}\rho \neq u_m + c$. $\square$

Proof of Theorem 4. The proof of this theorem is quite similar to the proof of Theorem 3, so we left it to the reader. $\square$

ACKNOWLEDGMENT

We would like to thanks A. Lopes, J. Rivera-Letelier and A. Tahzibi for the useful comments.

REFERENCES

- [1] de Lima, A. *Cohomologia e propriedades estocásticas de transformações expansoras e observáveis lipschitzianos*. Master's Thesis, ICMC-USP, 2007.
- [2] Baladi, V., *Positive Transfer Operators and Decay of Correlations*, Advanced Series in Non-linear Dynamics, vol. 16, World Scientific, (2000).
- [3] Bamón, R., Rivera-Letelier, J., Kiwi, J. and Urza, R., *On the Topology of Solenoidal Attractors of the Cylinder*, Ann. Inst. H. Poincaré Anal. Non Linéaire 23 (2006), no. 2, 209–236.
- [4] Broise, A. Transformations Dilatantes de L'Intervalle et Théorème Limites. in *Études Spectrales D'Opérateurs de Transfert et Applications*, Asterisque 238, (1996).
- [5] Viana, M., *Stochastic dynamics of deterministic systems*, Colóquio Brasileiro de Matemática-IMPA (1997).
- [6] Wong, S., *Holder Continuous Derivatives and Ergodic Theory*, J. of the London Mathematical Society s2-22(3), 506–520, 1980.

DEPARTAMENTO DE MATEMÁTICA, ICMC-USP, CAIXA POSTAL 668, SÃO CARLOS-SP, CEP 13560-970 SÃO CARLOS-SP, BRAZIL

E-mail address: `smania@icmc.usp.br`

URL: `www.icmc.usp.br/~smania/`